

anti- HSP70 antibody

Product Information

Catalog No.:	FNab04048
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3 , -20°C for 24 months (Avoid repeated freeze / thaw cycles.)

Background

HSPA1A, collectively known as HSP70(also referred to HSP72), is a stress-inducible member of heat-shock protein 70(HSP70) proteins which are highly conserved chaperons implicated in protein folding, protein refolding, protein transport, and protein targeting. Encoded by two closely linked, intronless and stress-inducible genes, HSPA1A and HSPA1B differ by only two amino acids and are believed to be fully interchangeable proteins. HSPA1A is a cytosol/nuclear protein able to translocate between cytoplasm and nucleus. Generally, HSPA1A is thought to be expressed in unstressed normal cells at low or undetectable levels. Expression of HSPA1A protein can be highly activated by various stressful stimuli. Significant up-regulation of HSPA1A has been found in various tumors. Recently it has been reported that HSPA1A can be constitutively expressed in selected cell types. HSP70 is a useful marker for exosome. This antibody well recognized the endogenous HSP70 protein in multiple cell lines.

Immunogen information

Immunogen:	heat shock 70kDa protein 1A
Synonyms:	Heat shock 70 kDa protein 1A,HSPA1A,Heat shock 70 kDa protein 1,HSP70-1,HSPA1, HSX70
Observed MW:	66-70 kDa
Uniprot ID :	PODMV8

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889


www.fn-test.com

Reactivity: Human, Mouse, Rat


Tested Application: ELISA, IHC, WB, IF, FC, IP

Recommended dilution: WB: 1:1000-1:10000; IP: 1:500-1:2000; IHC: 1:50-1:500; IF: 1:20-1:200


Image:




Immunohistochemistry of paraffin-embedded human liver cancer tissue slide using FNab04048(HSP70 Antibody) at dilution of 1:200



Immunofluorescent analysis of HepG2 cells, using HSPA1A antibody FNab04048 at 1:50 dilution. Blue pseudocolor = DAPI (fluorescent DNA dye).



IP Result of anti-HSP70 (IP: FNab04048, 3ug;
Detection: FNab04048 1:1000) with mouse brain
tissue lysate 3600ug.



HeLa cells were subjected to SDS PAGE followed
by western blot with FNab04048(HSP70
Antibody) at dilution of 1:5000